

# Vitamin D Level Status and hypertension among Elderly Iraqi People in Al Hillah City

Ismael Hasan Jawad<sup>1</sup>, Hasan ALwan Baiee<sup>2</sup>

<sup>1</sup>Student, PhD. Nursing Student, University of Babylon College of Nursing, <sup>2</sup>Prof. Dr. University of Babylon College of Nursing

## Abstract

**Background:** Vitamin D has different biological actions in the body . Vitamin D has the pleiotropic effects in multiple organ systems, and vitamin D deficiency was suggested to be associated with high blood pressure according to previous reports. Several interventional studies have examined the effect of vitamin D supplementation on high blood pressure patients

**Objective :** to identify Vitamin D level and its correlate with hypertension among old adult in Hilla city – Babylon province

## Introduction

Vitamin D, 25-hydroxyvitamin D (25(OH) D), is a dynamic fat-soluble vitamin that regulates calcium homeostasis and is important for bone and muscle health in people of all ages <sup>1</sup>. Vitamin D is logically present in some nutrients and dietary supplements and is formed endogenously when sunlight strikes the skin and motivates vitamin D synthesis. Serum concentration of 25-hydroxyvitamin D is the best indicator of vitamin D status in persons, with values of less than 30nmol/L (nmol/L = 0.4 ng/mL) measured to be insufficient for the universal health and wellbeing of adults <sup>2</sup>. A plethora of epidemiological and observational studies have established the correlation between vitamin D and general human wellbeing <sup>(3-5)</sup>. Studies suggest that adequate serum vitamin D of more than 30 nmol/L is concerned in avoiding cardiovascular disease <sup>6</sup> . An sufficient vitamin D serum level has also been informed to improve the immune system, avoid cancer, and limit its development <sup>7</sup>. Vitamin D is an essential part of nutrition. In association to the other vitamins, vitamin D has single metabolic and physiological special

effects<sup>(3-7)</sup> The deficiency of vitamin D is epidemically prevalent in the world; 20–25% of the population suffers from the deficiency of vitamin D in USA; Canada; Europe; Mexico; Asia; and Australia. <sup>8</sup> Surprisingly, the deficiency of vitamin D in the Persian Gulf nations is highly widespread, though there is adequate sunshine. The commonness of vitamin D (serum level of vitamin D) deficiency is upper among female adolescents and elderly in Iran and >80% in Saudi Arabia. Hypertension, also known as raised blood pressure, is a very common chronic disease and considered as a silent killer because it rarely causes symptoms<sup>(3)</sup>. Generally, older age, lower incomes and higher body mass index are proposed as the associated factors with the risk of hypertension<sup>46)</sup>. Accordingly, people having high blood pressure would increase in the condition of population ageing and prevalent westernized diet in Korean society<sup>(16)</sup>. Therefore, it is very important to look into the evidences and results about vitamin D in regards with its roles in controlling blood pressure at this point.

## Methodology

This was across sectional descriptive observational study included a non-probability ( convenient sample ) of elderly in Hilla City.

– Babylon province, the period of the study started from the first of January through August 2019, a pretested

---

### Corresponded Author:

Hasan ALwan Baiee,

University of Babylon College of Nursing

E-mail: hassanbaey@yahoo.com

questionnaire was used to interview the participants after obtaining their verbal consents, the sample included old adult, serum level of Vitamin D that made by chemo immunoassay method (maglumi instrument),The data were analyzed statistically to assess the associations between variables.

**Results**

The study included 300 participants 83.0% of the study sample had either insufficiency or deficiency of Vitamin D level. The proportion of adults with hypertension were more common among participants with deficient and insufficient vitamin D, 42% and 7% respectively , the difference was significant  $p < 0.05$ .

**Table (1) the mean age and number of participations.**


Parameter	Number	Mean of the age
65-69 years	168	67 years
70 -74 years	84	73 years
75-80and more	48	78 years
Total	300	

This table shows distribution of study participants according to the mean age and number of participations.

**Table (2) Means of vitamin D level by gender.**

Female	19.5% ng/ml
Male	25.8% ng/ml
Male and female	22.5% ng/ml

**This table shows distribution of study participants according to the means of vitamin D level by gender.**




**Figure (2) Means of vitamin D level by gender.**

**Table (3) Frequency distribution of the study group of age.**

Parameter	Frequency	percentage
65-69	168	56.0%
70 -74	84	28.0%
75-80 and more	48	16.0%
Total	300	100%

**This table shows distribution of study participants according to the age.**



**Figure (2) Frequency distribution of the study group by age.**

**Table (4) Association between Vitamin D Hypertension) among males.**

Study variables	Vitamin D level among male					$\chi^2$	P-value
	Deficiency (0-20 ng/ml)	Insufficiency (21-29 ng/ml)	Normal (30-100 ng/ml)	Total	Deficiency		
Hypertension Yes	37 (54.4%)	5 (26.3%)	2 (15.3%)	44	84.0%	57.99	< 0.002*
No	31 (45.6%)	14 (73.7%)	11 (84.7%)	56	55.3%		
Total	68 (100.0%)	19 (100.0%)	13 (100.0%)	99			

This Table shows Association between Vitamin D Hypertension) among males.

**Table (5) Association between Vitamin D and Hypertension)among females.**

Study variables	Vitamin D level among female					$\chi^2$	P-value
	Deficiency (0-20 ng/ml)	Insufficiency (21-29 ng/ml)	Normal (30-100 ng/ml)	Total	Deficiency		
Hypertension Yes	90 (66.1%)	16 (72.7%)	5 (11.6%)	111	81.0%	59.11	< 0.001*
No	46 (33.9%)	6 (27.3%)	38 (88.4%)	90	51.1%		
Total	136 (100.0%)	22 (100.0%)	43 (100.0%)	201			

This table shows Association between Vitamin D and Hypertension)among females.

## Discussion

In this study, the Vitamin D Level Status Among Iraqi Patients in Al Hillah City and the relation of these dependent variables to various epidemiological factors were assessed in 300 patients the most of sample are female 67% . The prevalence of vitamin D deficiency in those countries is higher in women than in men It seems that the skin complexion, poor sun exposure, vegetarian food habits and lack of vitamin D food fortification .

In this study most of patients are in the age group 65-69 years this result agree with (Heshmat et al 2008) who found that It was also indicated that vitamin D shortage is highest among individuals who are elderly, institutionalized, or hospitalized. It is recounted that 60% of the old adult in nursing homes were vitamin D deficient in the United States (Elliott et al.2003).

Findings of relationship between level of vitamin D and hypertension for male and female sample male and female have deficiency in the level of vitamin D (84 %) (81%). This results are agree with (Songcang 2017) who found that vitamin D deficiency is highest among people who have hypertension .

## Conclusion

Results of the current study showed a widespread, severe Vitamin D deficiency among participants of both sexes and in elderly people , urgent large scale public educational campaigns are needed to address this high priority public health problem in our society. We suggest that physicians should keep a check on the Vitamin D levels of elderly people in order to curb the ever-increasing incidence of hypertension.

**Financial Disclosure:** There is no financial disclosure.

**Conflict of Interest:** None to declare.

**Ethical Clearance:** All experimental protocols were approved under the University of Babylon College of Nursing, Iraq and all experiments were carried out in accordance with approved guidelines.

## References

1. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266-81.

2. Esteghamati A, Gouya MM, Abbasi M, Delavari A, Alikhani S, Alaedini F, *et al.* Prevalence of diabetes and impaired fasting glucose in the adult population of Iran: National survey of risk factors for non-communicable diseases of Iran. *Diabetes Care* 2008;31:96-8.
  3. Bringhurst FR, Demay MB, Krane SM, Kromenberg HM. Bone and mineral metabolism in health and disease. IKasper DL, Fauci AS, Longo DL, Braunwald E, Hauser SL, Jameson JL, editors. *Harrison's principles of internal medicine*. 16th ed. New York: McGraw Hill; 2005; 2238-48.
  4. Holick MF Sunlight , vitamin D. for bone health and prevention of autoimmune diseases, cancer, and cardiovascular disease. *Am J Clin Nutr*. 2004;80:1678S-88S. Age Pearson 1 0.038 0.221 0.315 0.102 0.080
  5. Marriott BM , Vitamin D. supplementation: A word of caution. *Ann Int Med* 1997;127:231-3.
  6. Norman AW. Sunlight, season, skin pigmentation, vitamin D, and 25-Hydroxyvitamin D: Integral components of the vitamin D endocrine system. *Am J Clin Nutr* 1998;67:1108-10.
  7. Compston JE, Vitamin D .deficiency: Time for action. *BMJ* 1998;317:1466-7
  8. Holick MF, Vitamin D ,deficiency *N Engl J Med* 2007;357:266-81.
  9. Mithal A, Wahl D, Bonjour JP, Burckhardt P, Dawson-Hughes B, Eisman J, *et al.* Global vitamin D status and determinants of hypovitaminosis D. *Osteoporosis Int.*, 2009;20:1807-20.
  10. Alvarez JA, Ashraf A. Role of vitamin d in insulin secretion and insulin sensitivity for glucose homeostasis. *Int J Endocrinol* 2010;2010:351-85.
- [1] National Institute of Health, Vitamin D. Fact Sheet for Health Professionals, May 2016, <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/>.
- [2] Institute of Medicine, Food and Nutrition Board, Dietary Reference Intakes for Calcium , Vitamin D, National Academy Press, Washington, DC, 2010.
- [3] MH Holick, Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis,” *The American Journal of Clinical Nutrition*, 2004;79( 3): 362–371.

- [4] A Hossein-nezhad, MF Holick, Vitamin D for health: a global perspective,” In *Mayo Clinic Proceedings*, 2013; 88 ( 7): 720–755.
- [5] MF Holick, Sunlight, vitamin D. for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease,” *The American Journal of Clinical Nutrition*, 2004;80( 6): 1678S–1688S.
- [6] I Mozos, O Marginean, “Links between vitamin D deficiency and cardiovascular diseases,” *BioMed Research International*, 2015;2015, Article ID 109275.
- [7] A Chirumbolo, “Vitamin D3 in cancer prevention and therapy: the nutritional issue,” *Hormone Molecular Biology and Clinical Investigation*, 2015; 23( 5): 71–78.
- [8] B Nikooyeh, T R Neyestani, M Farvid et al., “Daily consumption of vitamin D–or vitamin D+ calcium–fortified yogurt drink improved glycemic control in patients with type 2 diabetes: a randomized clinical trial,” *The American Journal of Clinical Nutrition*, 2011;93( 4): 764–771.
- [9] C Gagnon, Z X Lu, D J Magliano, et al., “Serum 25-hydroxyvitamin D, calcium intake, and risk of type 2 diabetes after 5 years: results from a national, population-based prospective study (the Australian Diabetes, Obesity and Lifestyle study),” *Diabetes Care*, 2011; 34( 5): 1133–1138.
- [10] RK Menon, A P Rickard, N Mannan, et al., “The effects of vitamin D2 or D3 supplementation on glycaemic control and related metabolic parameters in people at risk of type 2 diabetes: protocol of a randomised double-blind placebo-controlled trial,” *BMC Public Health*, 2013;13( 1): 999.
- [11] Bde Courten, A Mousa, N Naderpoor, H Teede, M de Courten, R cragg, “Vitamin D supplementation for the prevention of type 2 diabetes in overweight adults: study protocol for a randomized controlled trial,” *Trials*, 2015;16(1) : 335–346.
- [12] S Shab-Bidar, T R Neyestani, A Djazayeri, “Vitamin D receptor Cdx-2-dependent response of central obesity to vitamin D intake in the subjects with type 2 diabetes: a randomized clinical trial,” *The British Journal of Nutrition*, 2015;114(09): 1375–1384.
- [13] C Annweiler, G Allali, P Allain et al, “Vitamin D and cognitive performance in adults: a systematic review,” *European Journal of Neurology*, 2009;16( 10): 1083–1089,.
- [14] R E Anglin, Z Samaan, S D Walter, S D McDonald, “Vitamin D deficiency and depression in adults: systematic review and meta-analysis,” *The British Journal of Psychiatry*, 2013;202(2): 100–107.
- [15] American College of Cardiology. Serum 25-hydroxy Vitamin D in Ng/ml accepted. *American college of cardiology* 2011;58:1547-56..
16. Jo I, Ahn Y, Lee J, et al. Prevalence, awareness, treatment, control and risk factors of hypertension in Korea: the Ansan study. *J Hypertens*. 2001;19:1523–1532.
17. Lackland DT, Weber MA. Global burden of cardiovascular disease and stroke: hypertension at the core. *Can J Cardiol*. 2015;31:569–571.