

# Results of Surgical Treated of Nodular Thyroid in Women

**Rafa'a Sami Mahmood Al- Hayali**

*Specialist Surgeon, Fellow of Arab Board of Health Specialization, AL-jumhori Teaching Hospital ,  
Mosul, Iraq*

## Abstract

The long-term results of surgical treatment of nodular goiter and thyroid cancer were studied. Glands in 239 women under the age of 40 years. The patients were divided into two homogeneous groups: 1st — operated on for differentiated thyroid cancer; 2nd - operated on for benign neoplasms. Revealed causes and frequency the occurrence of recurrence of the disease, especially the course of postoperative hypothyroidism. The relationship of the occurrence of complications with the nosology and volume of the operation is analyzed. Evaluated the reproductive function of women after surgery. Tactics offered treatment of nodular goiter and thyroid cancer in pregnant women. Untreated thyroid diseases during pregnancy may lead to premature birth, preeclampsia (a severe increase in blood pressure), miscarriage, and low birth weight among other problems. Therefore, it is important to talk to your doctor if you have had a history of hypothyroidism or hyperthyroidism so you can be monitored before and during your pregnancy, and to be sure that your medication is properly adjusted, if necessary.

We studied the long-term results in 239 women under the age of 40 years. Patients were divided into two homogeneous groups: 1 - thyroid cancer, 2 - operated for benign tumors. The causes and frequency of recurrence of the disease and postoperative hypothyroidism were identified. The relation between the occurrence of complication's with nosology and operation volume was analyzed. We evaluated the reproductive function of treatment of the goiter and thyroid cancer in pregnant women.

**Keywords:** *Thyroidcancer, nodulargoiter, pregnant, treatment, postoperative*

## Introduction

The thyroid diseases—hyperthyroidism and hypothyroidism—are relatively common in pregnancy and important to treat. The thyroid is an organ located in the front of your neck that releases hormones that regulate your metabolism (the way your body uses energy), heart and nervous system, weight, body temperature, and many other processes in the body<sup>[1][2]</sup>.

Thyroid hormones are particularly necessary to assure healthy fetal development of the brain and nervous system during the first three months of your pregnancy since the baby depends on your hormones, which are delivered through the placenta. At around 12 weeks, the thyroid gland in the fetus will begin to produce its own thyroid hormones<sup>[3][4][5]</sup>.

There are 2 pregnancy-related hormones: estrogen and human chorionic gonadotropin (hCG) that may cause your thyroid levels to rise<sup>[6][7]</sup>. This may make it a bit harder to diagnose thyroid diseases that develop during pregnancy. However, your doctor will be on the look-out for symptoms that suggest the need for additional testing<sup>[8]</sup>.

However, if you have pre-existing hyperthyroidism or hypothyroidism, you should expect more medical attention to keep these conditions in control while you are pregnant, especially for the first trimester. Occasionally, pregnancy may cause symptoms similar to hyperthyroidism<sup>[9][10]</sup>; should you experience any uncomfortable or new symptoms, including palpitations, weight loss, or persistent vomiting, you should, of course, contact your physician<sup>[11]</sup>.

**Causes of Thyroid Disease in Pregnancy**  
Hyperthyroid disease—The most common cause of maternal hyperthyroidism during pregnancy is the autoimmune disorder Grave's disease. In this disorder, the body makes an antibody (a protein produced by the body when it thinks a virus or bacteria is present) called thyroid-stimulating immunoglobulin (TSI) that causes the thyroid to overreact and make too much thyroid hormone<sup>[12]</sup>.

Even if you've had radioactive iodine treatment or surgery to remove your thyroid, your body can still make the TSI antibody. If these levels rise too high, TSI will travel through your blood to the developing fetus, which may cause its thyroid to begin to produce more hormone than it needs<sup>[13]</sup>


<sup>[14]</sup>. So long as your doctor is checking your thyroid levels, both you and your baby will get the care needed to keep any problems in check<sup>[15]</sup>.

**Hypothyroid disease**—The most common cause of hypothyroidism is the autoimmune disorder known as Hashimoto's thyroiditis. In this condition, the body mistakenly attacks the cells of the thyroid gland, leaving the thyroid without enough cells and enzymes to make enough thyroid hormone to meet the body's needs<sup>[16][17]</sup>.

**Diagnosis of Thyroid Disease in Pregnancy**  
Hyperthyroidism and hypothyroidism in pregnancy are diagnosed based on symptoms, physical exam, and blood tests to measure levels of thyroid-stimulating hormone (TSH) and thyroid hormones T4, and for hyperthyroidism also T3<sup>[18]</sup>.

### **Thyroid physiology in pregnancy**

Approximately 94% of thyroid hormones are secreted by the thyroid gland as thyroxine or tetraiodothyronine (T4) and 6% as triiodothyronine (T3) (Fig. 1). T4 is catalytically converted to the more metabolically active T3 in peripheral tissues by deiodinases and a portion of peripherally-produced T3 returns to the circulation and it is because of this peripheral conversion that the plasma T4 to T3 ratio is approximately 4:1 <sup>[19]</sup> <sup>[20]</sup>. Both T4 and T3 are mostly bound to carrier proteins in the serum, chiefly thyroxine-binding globulin (TBG). However, it is the free hormones (free T4 (fT4) and free T3 (fT3)) that are available to be actively transported into cells and exert their effects<sup>[21][22]</sup>.



**Fig. 1 Hypothalamic-pituitary-thyroid axis and pregnancy.**

Changes in maternal thyroid function during pregnancy result from a combination of increased metabolic demands, increased serum TBG concentrations, stimulation of the TSH receptor by human chorionic gonadotropin (hCG) [23][24], an

increased mother-to-foetus transfer of thyroxine and an increased intraplacental breakdown of T4 and T3 (resulting from the placental expression of deiodinase 3)[25].

Total T4 and T3 concentrations increase by 50% as a result of a 50% increase in circulating TBG levels by 6–8 weeks of gestation; their levels plateau at around 16 weeks of gestation [26]. Maternal TSH is usually within normal limits during pregnancy but it can be decreased in the first trimester due to the increased hCG levels and the cross-reactivity of this hormone on TSH receptors<sup>[27]</sup> ; both are glycoprotein hormones with a common  $\alpha$  subunit and a considerable homology between their  $\beta$  subunits. Therefore hCG has a weak thyroid stimulating activity [28]. This hormonal interplay results in a biochemical picture of subclinical hyperthyroidism, which can

be considered as a physiological finding<sup>[29]</sup>. The decrease in hCG secretion later in pregnancy leads to reduction of serum fT4 and fT3 concentrations and finally the normalisation of TSH level<sup>[29]</sup>.

Thyroid hyperfunction and symptoms, if present, subside as hCG production falls, typically at 14–18 weeks of gestation. Ideally, the assay-specific TSH reference ranges for each trimester should be calculated based on the local population in iodine sufficient areas and pregnant women recruited for such calculations should be euthyroid and thyroid antibody negative. When this is not feasible, a reasonable alternative is to use the consensus ranges as per the various guidelines.

**Table 1: TSH reference ranges in pregnancy.**

	TSH reference ranges (mU/L)		
	First trimester	Second trimester	Third trimester
American Endocrine Society	0.1–2.5	0.2–3.0	0.3–3.0
American Thyroid Association			
European Thyroid Association	<2.5	<3.0	<3.5

Tasks:

1. Determination of risk factors recurrence of the disease after surgery.
2. Identification of the main causes of postoperative hypothyroidism and the possibility of its prevention.
3. Determine the degree of influence of these diseases and transferred operations on the state of

reproductive female systems.

## Materials and Methods

In AL-Gomhury General Hospital analyzed case histories of 239 women at the age of 40 years, which were operated on do nodal lesions of the thyroid gland for 4-year ago . They were divided into 2 groups. The first group consisted of 112 women operated on differentiated thyroid

cancer Shl. Isolation of various forms of thyroid cancer. PS based on the International Histological classification developed in 1974 by a group World

Health Organization Experts. The ratio of various forms of tumors is given in

**Table 1. Frequency of various forms of differentiated cancer thyroid gland**

Form of thyroid cancer	Number of patients	
	Abs	%
Papillary thyroid cancer	67	59.8
Follicular thyroid cancer	35	31.3
Follicular papillary thyroid	10	8.9
Total	112	100

The average age of patients was  $29.6 \pm 0.74$  years (from 10 up to 40 years).

In determining the stage of the process, the class TNM certification. By the time of surgery, the disease is was carried out in the following stages (Table 2).

The second group, consisting of 127 patients, were persons with benign thyroid pathology Shl. According to the results of postoperative histological remote material studies all observations were divided into groups (table 3).

**TABLE 2. The distribution of patients with thyroid cancer in stages**

Stage	Number of patients	
	Abs	Stage -%
T 1 N 0 M 0	16	Stage I - 14.3
T 2 N 0 M 0	58	
T 3 N 0 M 0	26	
T 2 N 1a M 0	3	Stage II - 77.7
T 4 N 0 M 0	5	
T 3 N 1a M 0	3	
T 3 N 1b M 0	1	
		Stage III - 8.0

**TABLE 3. Frequency of various forms of benign nodal lesions in patients of the 2nd group**

Nosological form	Number of patients	
	Abs	%
Colloid goiter	99	78
Thyroid adenomas	13	10.2
Autoimmune thyroiditis	15	11.8
Total	127	100

The average age of patients was  $30.6 \pm 0.6$  years (from 16 up to 40 years).

The following types of surgical procedures were performed for the patients. interventions (table 4).

**TABLE 4. Surgical interventions in patients of the 1st and 2nd groups**

Name of operation *	Group 1		Group 2	
	Abs	%	Abs	%
Thyroid lobe resection	0	0	18	14.2
Hemithyroidectomy	59	52.6	54	42.5
Thyroid lobe resection	1	0.9	29	22.8
Subtotal resection of the thyroid lobes	0	0	17	13.4
Subtotal resection of the thyroid gland	49	43.8	9	7.1
Thyroidectomy	3	2.7	0	0
Total	112	100	127	100

Note: \* - operation names are in accordance with nomenclature [10]

Thyroidectomy were performed at stages T 4 N 0 M 0 , T 3 N 0 M 0 and T 3 N 1b M 0 . In the latter case, thyroidectomy supplemented by bilateral fascial-dissecting tie fiber of the neck. Unilateral lymphadenectomy was performed in 6 cases (3 in addition to the subtotal resection of the thyroid gland and hemithyroidectomy) in the presence

of unilateral damage to regional lymph nodes. 4 patients with thyroid surgery combined with other goy operation (1 cholecystectomy by the traditional method, 2 laparoscopic cholecystectomy, 1 hernia repair with Mayo plastic surgery for umbilical hernia).

The early postoperative period was relatively satisfactorily in most patients. One patient (0.42%) developed bleeding, consuming battered surgical hemostasis. The phenomena of paresis of the gort not recorded in 4 patients (1.67%). In 2 cases required the imposition of a temporary tracheostoma (0.84%).

To study long-term results survey was carried out with the consent of the patient and included an tirovaniye (100%), survey (100%), ultrasonography of a thyroid gland zy (100%), reflexometry (100%), fine-needle asp biopsy (9.6%), a study of hormonal blood profile (84.9%), ECG (15%), examination by a gynecologistwith an assessment of the fertility function (4.6%). Tirescin- tigraphy despite its importance in identifying recurrence of thyroid cancer [11], according to technical reasons could be performed only in 5% of patients.

The survey questionnaire was asked questions about health status before and after surgery, family history of thyroid disease was studied glands, oncological diseases. Special attention

paid to childbearing function, quantity and course pregnancies, their connection with the operation and the method of completion.

For reflexometry, ref- Lexograph “Achilles-001”, developed by engineering technical cooperative “Novator” Omsk. For registration The result of the stratum was an electrocardiograph.

If a thyroid dysfunction is suspected glands produced determination of thyroid hormones new in the blood. For this outpatient on an empty stomach in patients determined the serum concentration of the following hormones: triiodothyronine (T 3 ), total thyroxin (T 4 ) and thyreotropin (TSH). Blood sampling was done before 11 o’clock in the morning from the cubital vein. Serum defended, frozen lived and kept at -20 ° C.

Ultrasound examination of the thyroid gland all 239 women of both groups were plagued. To conduct research was used apparatus Alloca, it’s often- that is 3.5 MHz.

**Table 5): Results and its discussion The frequency of recurrence of thyroid disease the gland was 2.7% for thyroid cancer and 9.4% for benign nodal thyropathies.**

Nosology	1st group	2nd group
Colloid goiter	-	9
Thyroiditis	-	3
Crayfish	3	-
Total	3 (2.7%)	12 (9.4%)

In this case, the re-formation of nodes was fixed 54 cases (22.6%), but their histological the structure differed from that established on the first operation (pseudorecidivy). Repeated operation was 21 patients were infected (8.8%). Higher percentage true recurrence in benign pathology was due to involvement in the pathological process a process of greater thyroid tissue and inadequate surgical intervention.

The relationship of the transaction volume with the frequency the recurrence of the disease. The following laws have been identified dimensionality:

- in case of benign pathology, relapse after subtotal resection of the thyroid glands are significantly lower than after hemithyroidectomy ( $p = 0.05$ );
- recurrence rate after hemithyroidectomy significantly higher in benign thyroid patty than in thyroid cancer ( $p = 0.05$ ).

Another important indicator for assessing long-term results of surgical treatment, is the presence of postoperative hypothyroidism.

When using the standardization method is established that if the patients of the 1st and 2nd groups were performed alone naive in terms of surgery, the frequency of hypothyroidism in the 2nd the group would be significantly higher ( $p = 0.05$ ).

Of the 34 women of the 1st group, hypothyroidism had 15 (44.1%), out of 30 of the 2nd group - 14 (46.7%). With a statistical treatment (Fig. 2) revealed that in the presence of hypothyroidism premature abortion rate significantly higher than that in the total populations, and in women without hypothyroidism ( $p = 0.01$ ). It also draws attention

to the fact that pregnancy is significantly less common absence of hypothyroidism than in the general population oscillations ( $p = 0.001$ ). This fact is indirectly about the high role of thyroid hormones in providing the normal course of pregnancy. Miscarriage of pregnancy with manifest hypothyroidism is more common than subclinical however, statistically this difference turned out to be true ( $p = 0.3$ ).

Hypothyroidism, present in 29 women during the menstruation and confirmed at that time level data TSH and T4, was subsequently compensated by the reception L-thyroxine in 17 patients. 12 women when receiving more than 200 mcg of L-thyroxine per day to achieve normalization of nya thyroid hormone fails. They were all previously surgical interventions were made in the amount of hemi- thyroidectomy and more.

### Findings

1. Hemithyroidectomy provides a low risk of re- diva only with differentiated thyroid cancer glands. With benign thyroid disease glands to reduce the likelihood of recurrence can perform subtotal resection of the thyroid gland PS or thyroidectomy.

2. The likelihood of postoperative hypothyroidism it depends not only on nosology and to a significant extent fines ( $\rho = -0.77 \pm 0.18$ ) is associated with the volume of the remaining tissue thyroid gland.

3. Violation of the reproductive function of women operated on for nodular lesions of the thyroid gland, is associated with the presence of postoperative hypothyroidism and does not depend on the morphological structure node tours.



**Conflict of interest:** There is no conflict of interest among the authors.

**Funding:** Self

**Ethical Clearance:** This study is ethically approved by the Institutional ethical Committee.

### References

1. Huscher CS. Endoscopic right thyroid lobectomy. *Surg Endosc.* 1997;11:877.
2. Miccoli P, Elisei R, Materazzi G, Capezzone M, Galleri D, Pacini F, Berti P, Pinchera A. Minimally invasive video-assisted thyroidectomy for papillary carcinoma: a prospective study of its completeness. *Surgery.* 2002 Dec 1;132(6):1070-4.
3. Chung YS, Choe JH, Kang KH, Kim SW, Chung KW, Park KS, Han W, Noh DY, Oh SK, Youn YK. Endoscopic thyroidectomy for thyroid malignancies: comparison with conventional open thyroidectomy. *World journal of surgery.* 2007 Dec;31(12):2302-6.
4. Jeong JJ, Kang SW, Yun JS, Sung TY, Lee SC, Lee YS, Nam KH, Chang HS, Chung WY, Park CS. Comparative study of endoscopic thyroidectomy versus conventional open thyroidectomy in papillary thyroid microcarcinoma (PTMC) patients. *Journal of surgical oncology.* 2009 Nov 1;100(6):477-80.
5. Lee KE, Rao J, Youn YK. Endoscopic thyroidectomy with the da Vinci robot system using the bilateral axillary breast approach (BABA) technique: our initial experience. *Surgical Laparoscopy Endoscopy & Percutaneous Techniques.* 2009 Jun 1;19(3):e71-5.
6. Kang S-W, Jeong JJ, Yun J-S, Sung TY, Lee SC, Lee YS, et al. Robot-assisted endoscopic surgery for thyroid cancer: experience with the first 100 patients. *Surg Endosc.* 2009;23:2399–406.
7. Kim WW, Jung JH, Park HY. A single surgeon's experience and surgical outcomes of 300 robotic thyroid surgeries using a bilateral axillo-breast approach. *J Surg Oncol.* 2015;111:135–40.
8. Noureldine SI, Abdelghani R, Saeed A, Cortes N, Abbas A, Aslam R, et al. Is robotic hemithyroidectomy comparable to its conventional counterpart? *Surgery.* 2013;154:363.
9. Yi O, Yoon JH, Lee Y-M, Sung T-Y, Chung K-W, Kim TY, et al. Technical and oncologic safety of robotic thyroid surgery. *Ann Surg Oncol.* 2013;20:1927–33.
10. -Song CM, Ji YB, Bang HS, Park CW, Kim DS, Tae K. Quality of life after robotic thyroidectomy by a gasless unilateral axillary approach. *Ann. Surg. Oncol.* 2014;21:4188-94.
- 11- Kim BS, Kang KH, Kang H, Park SJ. Central neck dissection using a bilateral axillo-breast approach for robotic thyroidectomy: comparison with conventional open procedure after propensity score matching. *Surg Laparosc Endosc Percutan Tech.* 2014;24:67–72.
- 12- Son SK, Kim JH, Bae JS, Lee SH. Surgical safety and oncologic effectiveness in robotic versus conventional open thyroidectomy in thyroid cancer: a systematic review and meta-analysis. *Ann Surg Oncol.* 2015;22:3022–32.
- 13- Tae K, Ji YB, Cho SH, Lee SH, Kim DS, Kim TW. Early surgical outcomes of robotic thyroidectomy by a gasless unilateral axillo-breast or axillary approach for papillary thyroid carcinoma: 2 years' experience. *Head Neck.*

- 2012;34:617–25.
- 14- Lee S, Ryu HR, Park JH, Kim KH, Kang S-W, Jeong JJ, et al. Early surgical outcomes comparison between robotic and conventional open thyroid surgery for papillary thyroid microcarcinoma. *Surgery*. 2012;151:724–30.
  - 15- Kim WW, Kim JS, Hur SM, Kim SH, Lee S-K, Choi JH, et al. Is robotic surgery superior to endoscopic and open surgeries in thyroid cancer? *World J Surg*. 2011;35:779–84.
  - 16- Tae K, Song CM, Ji YB, Kim KR, Kim JY, Choi YY. Comparison of surgical completeness between robotic total thyroidectomy versus open thyroidectomy. *Laryngoscope*. 2014;124:1042–7.
  - 17- Lang BH-H, Wong CKH, Tsang JS, Wong KP, Wan KY. A systematic review and meta-analysis comparing surgically-related complications between robotic-assisted thyroidectomy and conventional open thyroidectomy. *Ann Surg Oncol*. 2014;21:850–61
  - 18- Lee SG, Lee J, Kim MJ, Choi JB, Kim TH, Ban EJ, et al. Long-term oncologic outcome of robotic versus open total thyroidectomy in PTC: a case-matched retrospective study. *Surg Endosc*. 2015
  - 19- Kandil E, Hammad AY, Walvekar RR, Hu T, Masoodi H, Mohamed SE, et al. Robotic thyroidectomy versus nonrobotic approaches: a meta-analysis examining surgical outcomes. *Surg Innov*. 2016;23:317–25.
  - 20- Lee J, Nah KY, Kim RM, Ahn YH, Soh E-Y, Chung WY. Differences in postoperative outcomes, function, and cosmesis: open versus robotic thyroidectomy. *Surg Endosc*. 2010;24:3186–94.
  - 21- Tae K, Kim KY, Yun BR, Ji YB, Park CW, Kim DS, et al. Functional voice and swallowing outcomes after robotic thyroidectomy by a gasless unilateral axillo-breast approach: comparison with open thyroidectomy. *Surg Endosc*. 2012;26:1871–7.
  - 22- Lee J, Na KY, Kim RM, Oh Y, Lee JH, Lee J, et al. Postoperative functional voice changes after conventional open or robotic thyroidectomy: a prospective trial. *Ann Surg Oncol*. 2012;19:2963–70.
  - 23- Austin PC. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivar Behav Res*. 2011;46:399–424.
  - 24- Edge S, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A. *AJCC Cancer Staging Manual*. 7th ed. New York: Springer-Verlag; 2010.
  - 25- American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer. Cooper DS, Doherty GM, Haugen BR, Hauger BR, Kloos RT, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid Off J Am Thyroid Assoc*. 2009;19:1167–214.
  - 26- Sawka AM, Orlov S, Gelberg J, Stork B, Dowar M, Shaytzig M, et al. Prognostic value of postsurgical stimulated thyroglobulin levels after initial radioactive iodine therapy in well-differentiated thyroid carcinoma. *Head Neck*. 2008;30:693–700.
  - 27- Ryu IS, Song CI, Choi S-H, Roh J-L, Nam SY, Kim SY. Lymph node ratio of the central compartment is a significant predictor for

locoregional recurrence after prophylactic central neck dissection in patients with thyroid papillary carcinoma. *Ann Surg Oncol.* 2014;21:277–83.

28- Min HS. N stage: controversies and recent issues. *J Korean Thyroid Assoc.* 2012;5:109.